


First observation of production of three massive gauge bosons




Philip
Chang

Rice HEP Seminar
June 23, 2020



Univ. of California
San Diego

First observation of production of three massive gauge bosons $V = W, Z$




Philip
Chang

Rice HEP Seminar
June 23, 2020



Univ. of California
San Diego

First observation of production of three massive gauge bosons $V = W, Z$



Philip
Chang

Rice HEP Seminar
June 23, 2020



Univ. of California
San Diego



Discovery of Higgs boson

July 4, 2012




Discovery advanced our knowledge of origin of mass in a major way

Quarks



Leptons


Forces



- Is it the only Higgs boson?
(or are there more?)
- Are multi-*bosons* interactions SM?
- Are there more states involved in electroweak symmetry breaking?
- Is the Higgs potential SM?


Many more to be studied on electroweak sector at the LHC

Quarks





“massive” bosons

Forces




Leptons



- Is it the only Higgs boson?
(or are there more?)
- Are multi-*bosons* interactions SM?
- Are there more states involved in electroweak symmetry breaking?
- Is the Higgs potential SM?

Many more to be studied on electroweak sector at the LHC


More work to be done in electroweak sector



- Is it the only Higgs boson? (or are there more?)
- Are multi-*bosons* interactions SM?
- Are there more states involved in electroweak symmetry breaking?
- Is the Higgs potential SM?

Many more to be studied on electroweak sector at the LHC

More work to be done in electroweak sector




- Is it the only Higgs boson? (or are there more?)
- Are multi-bosons interactions SM?
- Are there more states involved in electroweak symmetry breaking?
- Is the Higgs potential SM?

1978 (way) before top/W/Z/Higgs discovery
Chanowitz, Furman, Hinchliffe

F, W^\pm , Z and H become “sthenons” in the sense of Appelquist and Bjorken [4]: they couple strongly to one another^{†1} but weakly to non-sthenons (i.e., the light particles in the theory).

Many more to be studied on electroweak sector at the LHC

Multi-“X” processes are rare and “heavy”



N.B. $x\text{sec} \times \text{decay BR}$ can be even smaller

multi-“massive”-X particles
 $X = t, W, Z, H$


Below picobarn most SM processes are electroweak multi-X production

We need LHC's large and energetic pp collision data

because rare

because “heavy”


Chang
UCSD




Multiply by 1000 to get the number of events produced for a picobarn process

During Run 2, CMS recorded 150 fb^{-1} of which 137 fb^{-1} have been validated as good quality data useable for physics analysis

LHC's large data enables us to study rare EW multi-X processes




Triboson process has access to studying many multi-*boson* interactions




Triboson process has access to studying many multi-*boson* interactions

Physics of VVV production ($V = W, Z$)




Triboson process has access to studying many multi-*boson* interactions

Physics of VVV production ($V = W, Z$)



Triboson process has access to studying many multi-*boson* interactions


Physics of VVV production ($V = W, Z$)



Triboson process has access to studying many multi-*boson* interactions


- ATLAS searched for WWW in 8 TeV: 0.96σ (1.05σ) arXiv:1610.05088
- CMS searched for WWW in 13 TeV 36 fb^{-1} : 0.6σ (1.78σ) arXiv:1905.04246
- ATLAS searched for VVV in 13 TeV 80 fb^{-1} : 4.1σ (3.1σ) arXiv:1903.10415

VVV evidence




arXiv:1903.10415

Axion-like-particle triboson signature limit



arXiv:1905.04246

SMEFT Dim8 operator limit



arXiv:1610.05088

Both ATLAS and CMS have been searching for triboson processes and using them to test SM and constrain new physics

Decay of W, Z bosons

$$W \rightarrow e/\mu (\sim 20\%)$$

$$Z \rightarrow ee/\mu\mu (\sim 7\%)$$

We select leptons w/
transverse momentum
(P_T) of $> 25, 20, 10$


τ decays in the
detector

$$\tau \rightarrow e, \mu + 2\nu$$

or

$$\tau \rightarrow \text{hadrons} + \nu$$

We include e, μ from τ 's from W/Z
decays in the analysis
But they have quite soft P_T and do
not pass the P_T requirements



W's and Z's produced can be identified via electrons and muons



CMS detector measures leptons very well

Chang
UCSD



e/μ among the **best**
measured particles at CMS
by combining tracker,
calorimeter, and chambers
measurements


(1-2% resolution for well measured ones)



Excellent lepton reconstruction and simulation at CMS

We need to further
classify the origin


N.B. electrons and muons
have different effects
(muons are cleaner)



A diagram showing a cone with a red vector labeled v and a dashed vector labeled v .



Dubbed “fake lepton”




The more leptons produced the lower the rate (i.e. lower bkg.)


Useful to organize physics analyses by N leptons

1. Organize analyses by leptons (likely) from W / Z
 - N leptons in the event
 - Flavor of the leptons
2. Reject events with b jets
3. Additional background suppression through smart choices
4. Reliably estimate the size of residual backgrounds
5. Observe VVV!

Smart humans and
smart machines
(Both cut / BDT)




Production cross section decreases with more Z's



Viable final states have O(fb) or less cross sections


Production cross section decreases with more Z's



Difficult final states
w/ bkg rate of
several orders of
magnitude larger

Viable final states have O(fb) or less cross sections

Production cross section decreases with more Z's



Difficult final states
w/ bkg rate of
several orders of
magnitude larger

$ZZZ \rightarrow 6L$
($L = e, \mu$)

11 attobarn

(~1.5 events produced
at Run 2 of LHC)

Viable



Viable final states have O(fb) or less cross sections

Target “fully” leptonic final states to go after first observation

One
exception

	Same-sign	3 leptons	4 leptons	5 leptons	6 leptons
Signals	$W^\pm \rightarrow l^\pm \nu$	$W \rightarrow l \nu$	$W \rightarrow l \nu$	$W \rightarrow l \nu$	$Z \rightarrow ll$
	$W^\pm \rightarrow l^\pm \nu$	$W \rightarrow l \nu$	$W \rightarrow l \nu$	$Z \rightarrow ll$	$Z \rightarrow ll$
	$W^\mp \rightarrow qq$	$W \rightarrow l \nu$	$Z \rightarrow ll$	$Z \rightarrow ll$	$Z \rightarrow ll$
	~2.5k evt.	~700 evt.	~140 evt.	~15 evt.	~1.5 evt.

Different modes populate different N lepton bins

Some cross contamination between N lepton bins exists but is small

Backgrounds in each N lepton region

	Same-sign	3 leptons	4 leptons	5 leptons	6 leptons
Signals	$W^\pm \rightarrow l^\pm \nu$ $W^\pm \rightarrow l^\pm \nu$ $W^\mp \rightarrow qq$	$W \rightarrow l \nu$ $W \rightarrow l \nu$ $W \rightarrow l \nu$	$W \rightarrow l \nu$ $W \rightarrow l \nu$ $Z \rightarrow ll$	$W \rightarrow l \nu$ $Z \rightarrow ll$ $Z \rightarrow ll$	$Z \rightarrow ll$ $Z \rightarrow ll$ $Z \rightarrow ll$
Dominant Bkgs.	$WZ \rightarrow l^\pm \nu l^\pm \bar{\nu}$ ^{lost} $t\bar{t} \rightarrow bb + l + X$ \hookrightarrow fake l	$WZ \rightarrow l \nu ll$ $t\bar{t} \rightarrow bb + ll + X$ \hookrightarrow fake l	$ZZ \rightarrow ll ll$ $ttZ \rightarrow ll ll + bbX$	$ZZ \rightarrow ll ll$ $+ \text{fake lep}$	$ZZ \rightarrow ll ll$ $+ 2 \text{ fake lep}$

N.B. $WZ \rightarrow 3l \sim 100k$, $ZZ \rightarrow 4l \sim 10k$

Once separated by N leptons dominant bkg. source becomes apparent

Backgrounds in each N lepton region


	Same-sign	3 leptons	4 leptons	5 leptons	6 leptons
Signals	$W^\pm \rightarrow l^\pm \nu$ $W^\pm \rightarrow l^\pm \nu$ $W^\mp \rightarrow qq$	$W \rightarrow l \nu$ $W \rightarrow l \nu$ $W \rightarrow l \nu$	$W \rightarrow l \nu$ $W \rightarrow l \nu$ $Z \rightarrow ll$	$W \rightarrow l \nu$ $Z \rightarrow ll$ $Z \rightarrow ll$	$Z \rightarrow ll$ $Z \rightarrow ll$ $Z \rightarrow ll$
Dominant Bkgs.	$WZ \rightarrow l^\pm \nu l^\pm l^\mp$ ^{lost} $t\bar{t} \rightarrow bb + l + X$ \hookrightarrow fake l	WW v. Z		$ZZ \rightarrow ll ll$ $ttZ \rightarrow ll ll + bbX$	$ZZ \rightarrow ll ll$ $+ 2 \text{ fake lep}$

N.B. $WZ \rightarrow 3l \sim 100k$, $ZZ \rightarrow 4l \sim 10k$

Selection on flavor and b tag will further reduce bkgs.

Once separated by N leptons dominant bkg. source becomes apparent

dilepton invariant mass of $Z \rightarrow \ell\ell$ decay



If one selects $|m_{\ell\ell} - m_Z| > 10 \text{ GeV}$ of $ee/\mu\mu$ final state Z is reduced by **an order** of magnitude

If one selects $e\mu$ final state, Z is reduced by **2 orders** of magnitude (e, μ from τ are soft)

\Rightarrow ZZ suppressed in 4 leptons: $ee/\mu\mu + e\mu$
 WZ suppressed in $e^\pm \mu^\mp e^\pm$

↑
 0 “SFOS”
 (Zero same-flavor opposite sign pair)

i.e. “ ee ” or “ $\mu\mu$ ”

Z decays predominantly to $ee/\mu\mu \Rightarrow$ select away from $Z \rightarrow ee/\mu\mu$

Splitting signal regions by lepton flavors




	Same-sign 2 leptons	3 leptons	4 leptons	5 leptons	6 leptons
Signals	$W^\pm \rightarrow l^\pm \nu$	$W \rightarrow l \nu$	$W \rightarrow l \nu$	$W \rightarrow l \nu$	$Z \rightarrow ll$
	$W^\pm \rightarrow l^\pm \nu$	$W \rightarrow l \nu$	$W \rightarrow l \nu$	$Z \rightarrow ll$	$Z \rightarrow ll$
	$W^\mp \rightarrow qq$	$W \rightarrow l \nu$	$Z \rightarrow ll$	$Z \rightarrow ll$	$Z \rightarrow ll$
	Split by $ee/e\mu/\mu\mu$ N.B. μ is cleaner than e	Split by # of SFOS e.g. 0: $e^\pm \mu^\mp e^\pm$ 1: $e^\pm e^\mp \mu^\pm$ 2: $e^\pm e^\mp e^\pm$	tag $Z \rightarrow ll$ then split $WW \rightarrow ee/\mu\mu$ v. $WW \rightarrow e\mu$		Not enough statistics single bin
	3 categories*	3 categories	2 categories*	1 category	1 category
	* marked ones will be further split				

Each N lepton analyses are further split by flavors

4



1. Organize analyses by “clean” leptons (likely) from W / Z
 - N leptons in the event
 - Flavor of the leptons
2. Reject events with b jets
3. Additional background suppression through smart choices
4. Reliably estimate the size of residual backgrounds
5. Observe VVV!

Smart humans and
smart machines
(Both cut / BDT)



Rejecting events with b jets



EW processes generally do not come
with b jets \Rightarrow Require # of $b = 0$



Signals do not have b jets

Added benefit of rejecting events with b

EW processes generally do not come
with b jets \Rightarrow Require # of $b = 0$




Signals do not have b jets

~~4~~

3

1. Organize analyses by “clean” leptons (likely) from W / Z
 - N leptons in the event
 - Flavor of the leptons
2. Reject events with b jets
3. Additional background suppression through smart choices
4. Reliably estimate the size of residual backgrounds
5. Observe VVV

Smart humans and
smart machines
(Both cut / BDT)



same-sign selection

Variable	m_{jj} -in and m_{jj} -out	1j
Trigger	Signal triggers, tab. 3.2	
Signal leptons	Exactly 2 tight SS leptons with $p_T > 25$ GeV	
Additional leptons	No additional very loose lepton	
Isolated tracks	No additional isolated tracks	
Jets	≥ 2 jets	1 jet
b-tagging	no b-tagged jets and soft b-tag objects	
$m_{\ell\ell}$	> 20 GeV	
$m_{\ell\ell}$	$ m_{\ell\ell} - m_Z > 20$ GeV if $e^\pm e^\pm$	
p_T^{miss}	> 45 GeV	
m_{JJ} (leading jets)	< 500 GeV	—
$\Delta\eta_{JJ}$ (leading jets)	< 2.5	—
m_{jj} (closest ΔR)	$65 < m_{jj} < 95$ GeV or $ m_{jj} - 80 \text{ GeV} \geq 15$ GeV	—
$\Delta R_{\ell j}^{\text{min}}$	—	< 1.5
m_T^{max}	> 90 GeV if not $\mu^\pm \mu^\pm$	> 90 GeV

Three leptons selection

Variable	0 SFOS	1 and 2 SFOS
Trigger	Signal triggers, tab. 3.2	
Signal leptons	3 tight leptons with charge sum = $\pm 1e$ $p_T > 25/25/25$ GeV $p_T > 25/20/20$ GeV	
Additional leptons	No additional very loose lepton	
m_{SFOS}	$m_{\text{SFOS}} > 20$ GeV and $ m_{\text{SFOS}} - m_Z > 20$ GeV	
$m_{\ell\ell\ell}$	$ m_{\ell\ell\ell} - m_Z > 10$ GeV	
SF lepton mass	> 20 GeV	—
Dielectron mass	$ m_{ee} - m_Z > 20$ GeV	—
Jets	≤ 1 jet	0 jets
b-tagging	No b-tagged jets and soft b-tag objects	
$\Delta\phi(\vec{p}_T(\ell\ell\ell), \vec{p}_T^{\text{miss}})$	—	> 2.5
$p_T(\ell\ell\ell)$	—	> 50 GeV
m_T^{3rd} (1 SFOS) or m_T^{max} (2 SFOS)	—	> 90 GeV

Four leptons selection

Variable	$e\mu$ category	$ee/\mu\mu$ category
Preselection	Selections in Table 20	
W candidate lepton flavors	$e\mu$	$ee/\mu\mu$
$m_{\ell\ell}$	Separated into 4 bins in (0, 40, 60, 100, ∞)	$ m_{\ell\ell} - m_Z > 10$ GeV
m_{T2}	$m_{T2} > 25$ GeV (for $m_{\ell\ell} > 100$ GeV)	...
$p_{T,4\ell}$ and p_T^{miss}	...	No $p_{T,4\ell}$ cuts and $p_T^{\text{miss}} > 120$ GeV (Bin A) $p_{T,4\ell} > 70$ GeV and $70 < p_T^{\text{miss}} < 120$ GeV (Bin B) $40 < p_{T,4\ell} < 70$ GeV and $70 < p_T^{\text{miss}} < 120$ GeV (Bin C)

5/6L will be
explained later

This is the full selections but I will not go in details for every single one

same-sign selection

Variable	m_{jj} -in and m_{jj} -out	1j
Trigger	Signal triggers, tab. 3.2	
Signal leptons	Exactly 2 tight SS leptons with $p_T > 25$ GeV	
Additional leptons	No additional very loose lepton	
Isolated tracks	No additional isolated tracks	
Jets	≥ 2 jets	1 jet

Split by N leptons
and requiring “Tight” leptons

$\Delta\eta_{jj}$ (leading jets)	< 2.5	—
m_{jj} (closest ΔR)	$65 < m_{jj} < 95$ GeV or $ m_{jj} - 80 \text{ GeV} \geq 15$ GeV	—
$\Delta R_{\ell j}^{\min}$	—	< 1.5
m_T^{\max}	> 90 GeV if not $\mu^\pm \mu^\pm$	> 90 GeV

Three leptons selection

Variable	0 SFOS	1 and 2 SFOS
Trigger	Signal triggers, tab. 3.2	
Signal leptons	3 tight leptons with charge sum = $\pm 1e$ $p_T > 25/25/25$ GeV $p_T > 25/20/20$ GeV	
Additional leptons	No additional very loose lepton	
m_{SFOS}	$m_{\text{SFOS}} > 20$ GeV and $ m_{\text{SFOS}} - m_Z > 20$ GeV	
$m_{\ell\ell\ell}$	$ m_{\ell\ell\ell} - m_Z > 10$ GeV	
SF lepton mass	—	
Dielectron mass	$ m_{ee} - m_Z > 20$ GeV	
Jets	≤ 1 jet	0 jets
b-tagging	No b-tagged jets and soft b-tag objects	
$\Delta\phi(\vec{p}_T(\ell\ell\ell), \vec{p}_T^{\text{miss}})$	—	> 2.5
$p_T(\ell\ell\ell)$	—	> 50 GeV
m_T^{3rd} (1 SFOS) or m_T^{\max} (2 SFOS)	—	> 90 GeV

Split by channels

Four leptons selection

Variable	$e\mu$ category	$ee/\mu\mu$ category
Preselection	Selections in Table 20	
W candidate lepton flavors	$e\mu$	$ee/\mu\mu$
$m_{\ell\ell}$	Separated into 4 bins in (0, 40, 60, 100, ∞)	$ m_{\ell\ell} - m_Z > 10$ GeV
m_{T2}	$m_{T2} > 25$ GeV (for $m_{\ell\ell} > 100$ GeV)	...
$p_{T,4\ell}$ and p_T^{miss}	...	No $p_{T,4\ell}$ cuts and $p_T^{\text{miss}} > 120$ GeV (Bin A) $p_{T,4\ell} > 70$ GeV and $70 < p_T^{\text{miss}} < 120$ GeV (Bin B) $40 < p_{T,4\ell} < 70$ GeV and $70 < p_T^{\text{miss}} < 120$ GeV (Bin C)

5/6L will be
explained later

But already you can notice a few things

same-sign selection

Variable	m_{jj} -in and m_{jj} -out	1j
Trigger	Signal triggers, tab. 3.2	
Signal leptons	Exactly 2 tight SS leptons with $p_T > 25$ GeV	
Additional leptons	No additional very loose lepton	
Isolated tracks	No additional isolated tracks	
Jets	≥ 2 jets	1 jet
b-tagging	no b-tagged jets and soft b-tag objects	
$m_{\ell\ell}$	> 20 GeV	
$m_{\ell\ell}$	$ m_{\ell\ell} - m_Z > 20$ GeV if $e^\pm e^\pm$	
p_T^{miss}	> 45 GeV	
m_{jj} (leading jets)	< 500 GeV	—
$\Delta\eta_{jj}$ (leading jets)	< 2.5	—
m_{jj} (closest ΔR)	$65 < m_{jj} < 95$ GeV or $ m_{jj} - 80 \text{ GeV} \geq 15$ GeV	—
$\Delta R_{\ell j}^{\text{min}}$	—	< 1.5
m_T^{max}	> 90 GeV if not $\mu^\pm \mu^\pm$	> 90 GeV

Three leptons selection

Variable	0 SFOS	1 and 2 SFOS
Trigger	Signal triggers, tab. 3.2	
Signal leptons	Exactly 3 tight SS leptons with $p_T > 25$ GeV	
Additional leptons	No additional very loose lepton	
m_{SFOS}	> 20 GeV	
$m_{\ell\ell\ell}$	> 20 GeV	
SF lepton mass	> 20 GeV	
Dielectron mass	> 20 GeV	
Jets	≥ 2 jets	
b-tagging	no b-tagged jets and soft b-tag objects	
$\Delta\phi(\vec{p}_T(\ell\ell\ell), \vec{p}_T^{\text{miss}})$	—	> 2.5
$p_T(\ell\ell\ell)$	—	> 50 GeV
m_T^{3rd} (1 SFOS) or m_T^{max} (2 SFOS)	—	> 90 GeV

- Split by number of jets
- Dijet invariant mass: m_{jj}
- Transverse mass: m_T
- “S”transverse mass: m_{T2}
- Missing transverse energy


Four leptons selection

Variable	$e\mu$ category	$ee/\mu\mu$ category
Preselection	Selections in Table 20	
W candidate lepton flavors	$e\mu$	$ee/\mu\mu$
$m_{\ell\ell}$	Separated into 4 bins in (0, 40, 60, 100, ∞)	$ m_{\ell\ell} - m_Z > 10$ GeV
m_{T2}	$m_{T2} > 25$ GeV (for $m_{\ell\ell} > 100$ GeV)	...
$p_{T,4\ell}$ and p_T^{miss}	...	No $p_{T,4\ell}$ cuts and $p_T^{\text{miss}} > 120$ GeV (Bin A) $p_{T,4\ell} > 70$ GeV and $70 < p_T^{\text{miss}} < 120$ GeV (Bin B) $40 < p_{T,4\ell} < 70$ GeV and $70 < p_T^{\text{miss}} < 120$ GeV (Bin C)


5/6L will be
explained later

But I will highlight these 5 points in the coming slides


Reconstruct $W \rightarrow qq$ in $WWW \rightarrow l\bar{l}qq$




N.B. some signals are outside the window
(See next slide)



dijet invariant mass for signal peaks around W mass



Difficult to match $W \rightarrow qq$
⇒ Select off-W-mass peak region



Difficult to reconstruct both jets
⇒ Select 1 jet (1J) events

2 additional categories (m_{jj} -in, m_{jj} -out, 1J) each split by $ee/e\mu/\mu\mu$
⇒ Total of 9 signal regions for same-sign analysis

We cover wide range of possible jet final states to maximize sensitivity

Kinematic endpoints for 3 leptons

Separated by # of SFOS pairs:

- 0 SFOS (low bkg.)
- 1 SFOS
- 2 SFOS

0SFOS is by far the cleanest

For 1SFOS it is clear which one is from W:

$$\frac{e^{\pm}e^{\mp}}{Z} \quad \frac{\mu^{\pm}}{W} \quad \frac{\mu^{\pm}\mu^{\mp}}{Z} \quad \frac{e^{\pm}}{W}$$



For 2SFOS it is less clear which one is from W:

$$\frac{e^{\pm}}{W?} \quad \frac{e^{\mp}}{W?} \quad \frac{e^{\pm}}{W?} \quad \frac{\mu^{\pm}}{W?} \quad \frac{\mu^{\mp}}{W?} \quad \frac{\mu^{\pm}}{W?}$$

Take max m_T computed from either leptons

⇒ 3 signal regions for 3 leptons


$$m_T^{3rd} (1 \text{ SFOS}) \text{ or } m_T^{\max} (2 \text{ SFOS}) > 90 \text{ GeV}$$



By flavor, W lepton can be identified and kinematic endpoints can be used

4 lepton event

CMS experiment at the LHC, CERN
Data recorded: 2016-Jul-23 08:13:27.898048 GMT
Run 277168, Event No. 3219714497 LS 1799



Kinematic endpoints for 4 leptons



Events are separated into 2 categories by flavor:

- “ $e\mu$ channel”: $(ee/\mu\mu)_{\text{on-Z-mass}} + e\mu$ (low bkg.)
- “ $ee/\mu\mu$ channel”: $(ee/\mu\mu)_{\text{on-Z-mass}} + ee/\mu\mu$

$e\mu$ channel utilizes m_{T2} variable, which is a generalization of m_T for multiple missing particles. m_{T2} is sensitive to the end points of m_τ from $ZZ \rightarrow \ell\ell\tau\tau$

ZZ bkg in $ee/\mu\mu$ have low missing energy

Combine these and a few more kinematic variables to form total of 7 signal regions for 4 lepton analysis



Exploit differences between $Z \rightarrow \ell\ell$ v. $WW \rightarrow \ell\nu\ell\nu$


5 leptons target $W \underline{Z} \underline{Z}$ signal

Require the 5 lepton events to contain two SFOS pair consistent with Z mass

The dominant background is $ZZ \rightarrow \ell\ell\ell\ell$ plus a fake lepton

The fake lepton has low transverse mass while the signal's W has transverse mass peaking at W mass

Cut-and-count of one bin



Exploit the features of $W \rightarrow \ell \nu$ decay


6 leptons

Select at least 6 leptons

Require $\Sigma P_T \geq 250$ GeV

Less than 1 event expected

Very clean channel



Not enough stats, so search inclusively


~~5~~ steps to VVV observation

~~4~~


3

1. Organize analyses by “clean” leptons (likely) from W / Z
 - N leptons in the event
 - Flavor of the leptons
2. Reject events with b jets
3. Additional background suppression through smart choices
4. Reliably estimate the size of residual backgrounds
5. Observe VVV!

Smart humans and
smart machines
(Both cut / **BDT**)



Boosted decision tree is widely used in many analyses at the LHC



https://arogozhnikov.github.io/2016/07/05/gradient_boosting_playground.html

Train dedicated boosted decision trees to maximize sensitivity

	Same-sign	3 leptons	4 leptons	5 leptons	6 leptons
Signals	$W^\pm \rightarrow l^\pm \nu$ $W^\pm \rightarrow l^\pm \nu$ $W^\mp \rightarrow qq$	$W \rightarrow l \nu$ $W \rightarrow l \nu$ $W \rightarrow l \nu$	$W \rightarrow l \nu$ $W \rightarrow l \nu$ $Z \rightarrow ll$	$W \rightarrow l \nu$ $Z \rightarrow ll$ $Z \rightarrow ll$	$Z \rightarrow ll$ $Z \rightarrow ll$ $Z \rightarrow ll$
Dominant Bkgs.	<div> $WZ \rightarrow l^\pm \nu l^\pm \bar{\nu}$ <p>lost \nearrow</p> </div> <div> $t\bar{t} \rightarrow bb + l + X$ \hookrightarrow fake l </div>	<div> $WZ \rightarrow l \nu ll$ </div> <div> $t\bar{t} \rightarrow bb + ll + X$ \hookrightarrow fake l </div>	<div> $ZZ \rightarrow ll ll$ </div> <div> $t\bar{t}Z \rightarrow ll ll + bbX$ </div>	$ZZ \rightarrow ll ll$ + fake lep	$ZZ \rightarrow ll ll$ + 2 fake lep

“Prompt” bkgs.



“Fake” bkgs.

$t\bar{t}Z$ bkg.

ZZ bkg.

No BDT trained for
5/6 leptons (not
enough stats)

Train different BDTs against different backgrounds




Maintained same categorizations but cut on BDT to maximize sensitivity


Total number of bins stayed same (9 for same-sign, 3 for 3 leptons)

Cut on each BDT scores to create a high sensitivity bin

2D plane in BDT scores for 4 lepton
 $Z \rightarrow \ell\ell + e\mu$ event category



5 bins are created
from 2D planes



**For $Z \rightarrow \ell\ell + ee/\mu\mu$ event category, 2 bins are created (not shown)

Created multiple bins in BDTs to maximize sensitivity

~~5~~ steps to VVV observation


~~4~~

~~3~~

2

1. Organize analyses by “clean” leptons (likely) from W / Z
 - N leptons in the event
 - Flavor of the leptons
2. Reject events with b jets
3. Additional background suppression through smart choices
4. Reliably estimate the size of residual backgrounds
5. Observe VVV!

Smart humans and
smart machines
(Both cut / BDT)





Now two steps left

	Same-sign 2 leptons	3 leptons	4 leptons	5 leptons	6 leptons
Dominant Bkgs.	$WZ \rightarrow l^\pm \nu l^\pm \bar{\nu}$ (lost ν) $t\bar{t} \rightarrow bb + l + X$ \hookrightarrow fake l	$WZ \rightarrow l \nu ll$ $t\bar{t} \rightarrow bb + ll + X$ \hookrightarrow fake l	$ZZ \rightarrow ll ll$ $t\bar{t}Z \rightarrow ll ll + bbX$	$ZZ \rightarrow ll ll$ $+ \text{fake lep}$	$ZZ \rightarrow ll ll$ $+ 2 \text{ fake lep}$

Types of backgrounds	Suppressed via	Bkg. estimation
Fake leptons	Isolation	Reliably extrapolate across isolation
Backgrounds with b jets	b tagging	Reliably extrapolate across b tagging
Lost leptons	Removing events with 3rd lepton	Reliably extrapolate across N leptons
Irreducible	Smart flavor choices	Reliably extrapolate across flavor

Reliably extrapolate across the method used to suppress background to estimate the size of residual backgrounds in signal region

Pick one lepton events with phase space dominated by QCD (dijet) events



Fake rate is then applied to signal like region with "Loose"-ly identified leptons

"Side band" in isolation

Underlying effects (P_T of quarks) that govern fake rate are not measurable
 \Rightarrow **Source of systematics (~30%)**

Estimate fake lepton by measuring fake rate from QCD events


Lepton finding efficiency is well modeled by MC

(factors: P_T , η , lepton ID)

Construct a control region with 3 leptons and
extrapolate across 3 lepton \rightarrow 2 leptons

Experimental systematics assigned

Control region data statistics dominates
uncertainty (20%)




Estimate lost lepton background by extrapolating across # of leptons

Backgrounds with b jets / irreducible


Devise control regions and extrapolate to signal region

N_b in 3 lepton




Extrapolate across N_b tag ($\sim 10\%$)

4 lepton BDT score
 $Z \rightarrow ll + e\mu + b$ jets



4 lepton m_{T2}
 $Z \rightarrow ll + ee/\mu\mu$



Extrapolate across flavor
(uncertainty $\sim 5\%$)

Extrapolate from control region to estimate backgrounds

Putting it all together

	Same-sign 2 leptons	3 leptons	4 leptons	5 leptons	6 leptons
Signals	$W^\pm \rightarrow l^\pm \nu$ $W^\pm \rightarrow l^\pm \nu$ $W^\mp \rightarrow qq$	$W \rightarrow l \nu$ $W \rightarrow l \nu$ $W \rightarrow l \nu$	$W \rightarrow l \nu$ $W \rightarrow l \nu$ $Z \rightarrow ll$	$W \rightarrow l \nu$ $Z \rightarrow ll$ $Z \rightarrow ll$	$Z \rightarrow ll$ $Z \rightarrow ll$ $Z \rightarrow ll$
Split Flavor	3	3	2	1	1
Channel specific splits	mjj-in mjj-out 1J	-	Split in kinematics or BDT	-	-
Total	9 bins	3 bins	7 bins	1 bin	1 bin

Total of 21 bins

~~5~~ steps to VVV observation

~~4~~

~~3~~

~~2~~


1

1. Organize analyses by “clean” leptons (likely) from W / Z
 - N leptons in the event
 - Flavor of the leptons
2. Reject events with b jets
3. Additional background suppression through smart choices
4. Reliably estimate the size of residual backgrounds
5. Observe VVV!

Let's observe!

Results (BDT-based analysis)


$$\text{Signal strength } \mu = \frac{\text{Measured cross section}}{\text{Theoretical cross section}}$$



BDT-based analysis is more sensitive so this is our main result


Results (Cut-based analysis)

$$\text{Signal strength } \mu = \frac{\text{Measured cross section}}{\text{Theoretical cross section}}$$



Cut-based analysis is also reported for cross check and completeness
(also easier to understand by theorists if re-interpreted)

VVV mode	Significance [σ]
WWW	3.3 (3.1)
WWZ	3.4 (4.1)
WZZ	1.7 (0.7)
ZZZ	0 (0.9)
Combined	5.7 (5.9)






$$\text{Signal strength } \mu = \frac{\text{Measured cross section}}{\text{Theoretical cross section}}$$

- We have **observed** production of three massive gauge boson for the **first** time!
- We also found **evidences** separately for the WWW and WWZ production.
- The cross sections are compatible with the standard model expectation.



First observation of VVV and evidences for WWW and WWZ productions

Now that we have established VVV production we can use it to test SM and also search new physics (cf. Four fermion interaction with Fermi constant)



Establishment of VVV production opens up a new physics program

Fully leptonic v. Semi leptonic channel




—— If BSM exists, effects are same ——



- Physics of $V \rightarrow f\bar{f}$ is well understood
- We have now established $pp \rightarrow VVV$ production in “fully” leptonic decay
- Therefore, there ought to be $pp \rightarrow VVV \rightarrow$ semi-leptonic
 \Rightarrow If new physics alters $pp \rightarrow VVV$, it will alter fully / semi leptonic the same

$VVV \rightarrow$ semi-leptonic ought to have same physics as $VVV \rightarrow$ fully leptonic


Fully leptonic v. Semi leptonic channel



Clean channel for discovery but probing tail is **difficult**



Bkg is larger but distinct high P_T feature can **discriminate** bkg.



Signal Bkg. Small Large Signal Bkg.


We can probe $VVV \rightarrow$ semi-leptonic for new physics

More multi-massive-X processes for future


listing a few multi-massive-X processes with **same-sign**

$$pp \rightarrow W^\pm W^\pm H$$


*Same-sign
is special*



$$pp \rightarrow t W^\pm W^\pm j$$




High P_T top
(> 500 GeV)




Many more

WWH, tWWj, ttWW, ttZH

$$pp \rightarrow ttWW$$



$$pp \rightarrow ttZH$$



There are many more multi-massive-X production to be explored at LHC


- First observation of VVV productions was made by CMS collaboration
- Also found evidences for WWW and WWZ
- first hints for WZZ production and no hints for ZZZ yet
- The measured cross section is compatible with SM
- This establishes VVV process and opens a unique opportunity to test SM
- New physics can be also searched
- In the future, LHC will continue to probe electroweak interactions in various VVV channel
- Also, LHC will have to continue to probe unexplored electroweak productions of massive particles



Backup



Compact Muon Solenoid LHC, CERN




Visit us: [CMS Public Website](#), [CMS Physics](#) ; Contact us: [CMS Publications Committee](#)

CMS Publications

1000	SMP-19-014	Observation of the production of three massive gauge bosons at $\sqrt{s} = 13$ TeV	Submitted to PRL	19 June 2020
999	HIN-19-001	Evidence for top quark production in nucleus-nucleus collisions	Submitted to NP	19 June 2020
998	TRG-17-001	Performance of the CMS Level-1 trigger in proton-proton collisions at $\sqrt{s} = 13$ TeV	Submitted to JINST	18 June 2020

April 2020

CMS Preliminary



Quantities	WWW	WWZ	WZZ	ZZZ
$\sigma_{pp \rightarrow VVV \text{ non-VH}} \text{ (fb)}$	216.0	165.1	55.7	14.0
$\sigma_{\text{VH} \rightarrow VVV} \text{ (fb)}$	293.4	188.9	36.0	23.1
$\sigma_{\text{total}} \text{ (fb)}$	509.4	354.0	91.6	37.1
$\mathcal{B}_{VVV \rightarrow SS} \text{ (%)}$	7.16	-	-	-
$\mathcal{B}_{VVV \rightarrow 3\ell} \text{ (%)}$	3.46	4.82	6.37	-
$\mathcal{B}_{VVV \rightarrow 4\ell} \text{ (%)}$	-	1.16	0.81	3.22
$\mathcal{B}_{VVV \rightarrow 5\ell} \text{ (%)}$	-	-	0.39	-
$\mathcal{B}_{VVV \rightarrow 6\ell} \text{ (%)}$	-	-	-	0.13
$\sigma_{\text{total}} \times \mathcal{B}_{VVV \rightarrow SS} \text{ (fb)}$	36.4	-	-	-
$\sigma_{\text{total}} \times \mathcal{B}_{VVV \rightarrow 3\ell} \text{ (fb)}$	17.6	17.1	5.83	-
$\sigma_{\text{total}} \times \mathcal{B}_{VVV \rightarrow 4\ell} \text{ (fb)}$	-	4.12	0.74	1.19
$\sigma_{\text{total}} \times \mathcal{B}_{VVV \rightarrow 5\ell} \text{ (fb)}$	-	-	0.36	-
$\sigma_{\text{total}} \times \mathcal{B}_{VVV \rightarrow 6\ell} \text{ (fb)}$	-	-	-	0.05
$\sigma_{\text{total}} \times \mathcal{B}_{VVV \rightarrow SS} \times 137 \text{ fb}^{-1} (N_{\text{evts}})$	4987	-	-	-
$\sigma_{\text{total}} \times \mathcal{B}_{VVV \rightarrow 3\ell} \times 137 \text{ fb}^{-1} (N_{\text{evts}})$	2411	2343	799	-
$\sigma_{\text{total}} \times \mathcal{B}_{VVV \rightarrow 4\ell} \times 137 \text{ fb}^{-1} (N_{\text{evts}})$	-	564	101	163
$\sigma_{\text{total}} \times \mathcal{B}_{VVV \rightarrow 5\ell} \times 137 \text{ fb}^{-1} (N_{\text{evts}})$	-	-	49.3	-
$\sigma_{\text{total}} \times \mathcal{B}_{VVV \rightarrow 6\ell} \times 137 \text{ fb}^{-1} (N_{\text{evts}})$	-	-	-	6.85

Features	Selections		
	SS + $\geq 2j$	SS + 1j	3ℓ
Triggers	Select events passing dilepton triggers		
Number of leptons	Select events with 2 (3) leptons passing SS-ID (3ℓ -ID) for SS (3ℓ) final states		
Number of leptons	Select events with 2 (3) leptons passing veto-ID for SS (3ℓ) final states		
Isolated tracks	No additional isolated tracks		—
b-tagging	no b-tagged jets and soft b-tag objects		
Jets	≥ 2 jets	1 jet	≤ 1 jet
m_{JJ} (leading jets)	< 500 GeV		—
$\Delta\eta_{JJ}$ (leading jets)	< 2.5		—
$m_{\ell\ell}$	> 20 GeV		—
$m_{\ell\ell}$	$ m_{\ell\ell} - m_Z > 20$ GeV if $e^\pm e^\pm$		—
m_{SFOS}	—	—	$m_{SFOS} > 20$ GeV
m_{SFOS}	—	—	$ m_{SFOS} - m_Z > 20$ GeV
$m_{\ell\ell\ell}$	—	—	$ m_{\ell\ell\ell} - m_Z > 10$ GeV


Variable	m_{jj} -in and m_{jj} -out	1j
Trigger	Signal triggers, tab. 3.2	
Signal leptons	Exactly 2 tight SS leptons with $p_T > 25$ GeV	
Additional leptons	No additional very loose lepton	
Isolated tracks	No additional isolated tracks	
Jets	≥ 2 jets	1 jet
b-tagging	no b-tagged jets and soft b-tag objects	
$m_{\ell\ell}$	> 20 GeV	
$m_{\ell\ell}$	$ m_{\ell\ell} - m_Z > 20$ GeV if $e^\pm e^\pm$	
p_T^{miss}	> 45 GeV	
m_{JJ} (leading jets)	< 500 GeV	—
$\Delta\eta_{JJ}$ (leading jets)	< 2.5	—
m_{jj} (closest ΔR)	$65 < m_{jj} < 95$ GeV or $ m_{jj} - 80 \text{ GeV} \geq 15$ GeV	—
$\Delta R_{\ell j}^{\text{min}}$	—	< 1.5
m_T^{max}	> 90 GeV if not $\mu^\pm \mu^\pm$	> 90 GeV

Variable	0 SFOS	1 and 2 SFOS
Trigger	Signal triggers, tab. 3.2	
Signal leptons	3 tight leptons with charge sum = $\pm 1e$ $p_T > 25/25/25$ GeV $p_T > 25/20/20$ GeV	
Additional leptons	No additional very loose lepton	
m_{SFOS}	$m_{\text{SFOS}} > 20$ GeV and $ m_{\text{SFOS}} - m_Z > 20$ GeV	
$m_{\ell\ell\ell}$	$ m_{\ell\ell\ell} - m_Z > 10$ GeV	
SF lepton mass	> 20 GeV	—
Dielectron mass	$ m_{ee} - m_Z > 20$ GeV	—
Jets	≤ 1 jet	0 jets
b-tagging	No b-tagged jets and soft b-tag objects	
$\Delta\phi(\vec{p}_T(\ell\ell\ell), \vec{p}_T^{\text{miss}})$	—	> 2.5
$p_T(\ell\ell\ell)$	—	> 50 GeV
m_T^{3rd} (1 SFOS) or m_T^{max} (2 SFOS)	—	> 90 GeV

Features	Selections
Number of leptons	Select events with 4 leptons passing common veto-ID
Triggers	Select events passing dilepton triggers
Z lepton	Find opposite charge lepton pairs, passing ZID, closest to m_Z Require Z leptons to have $p_T > 25, 15$ GeV
W lepton	Require that leftover leptons are opposite charge and pass WID Require W leptons to have $p_T > 25, 15$ GeV
Low mass resonances	Require any opposite charge pair invariant mass to be greater than 12 GeV
b-tagged jets	no b-tagged jet
Z mass window	Require invariant mass of the Z leptons to be within 10 GeV of Z boson mass


Variable	$e\mu$ category	$ee/\mu\mu$ category
Preselection	Selections in Table 20	
W candidate lepton flavors	$e\mu$	$ee/\mu\mu$
$m_{\ell\ell}$	Separated into 4 bins in (0, 40, 60, 100, ∞)	$ m_{\ell\ell} - m_Z > 10 \text{ GeV}$
m_{T2}	$m_{T2} > 25 \text{ GeV}$ (for $m_{\ell\ell} > 100 \text{ GeV}$)	...
$p_{T,4\ell}$ and p_T^{miss}	...	No $p_{T,4\ell}$ cuts and $p_T^{\text{miss}} > 120 \text{ GeV}$ (Bin A) $p_{T,4\ell} > 70 \text{ GeV}$ and $70 < p_T^{\text{miss}} < 120 \text{ GeV}$ (Bin B) $40 < p_{T,4\ell} < 70 \text{ GeV}$ and $70 < p_T^{\text{miss}} < 120 \text{ GeV}$ (Bin C)

$$m_{T2} = \min_{\vec{p}_T^{\nu(1)} + \vec{p}_T^{\nu(2)} = \vec{p}_T^{\text{miss}}} \left[\max \left(m_T^{(1)}(\vec{p}_T^{\nu(1)}, \vec{p}_T^e), m_T^{(2)}(\vec{p}_T^{\nu(2)}, \vec{p}_T^{\mu}) \right) \right]$$





For $WW \rightarrow l\nu l\nu$ sub-system of WWZ , endpoint is at m_W


For $Z \rightarrow \tau\tau \rightarrow ll\nu\nu\nu\nu$ sub-system of ZZ , endpoint is at m_τ





	ZZ BDT range	ttZ BDT range
$e\mu$ BDT bin 1	$(-\infty, -0.908)$	$(-\infty, \infty)$
$e\mu$ BDT bin 2	$(-0.908, \infty)$	$(-\infty, 0.015)$
$e\mu$ BDT bin 3	$(-0.908, 0.733)$	$(0.015, \infty)$
$e\mu$ BDT bin 4	$(0.733, \infty)$	$(0.015, 3.523)$
$e\mu$ BDT bin 5	$(0.733, \infty)$	$(3.523, \infty)$
$ee/\mu\mu$ BDT bin A	$(0, 3)$	-
$ee/\mu\mu$ BDT bin B	$(3, \infty)$	-











Process	Higgs boson contributions as signal		Higgs boson contributions as background	
	sequential-cut	BDT-based	sequential-cut	BDT-based
WWW	2.5 (2.9)	3.3 (3.1)	1.0 (1.8)	1.6 (1.9)
WWZ	3.5 (3.6)	3.4 (4.1)	0.9 (2.2)	1.3 (2.2)
WZZ	1.6 (0.7)	1.7 (0.7)	1.7 (0.8)	1.7 (0.8)
ZZZ	0.0 (0.9)	0.0 (0.9)	0.0 (0.9)	0.0 (0.9)
VVV	5.0 (5.4)	5.7 (5.9)	2.3 (3.5)	2.9 (3.5)

Process	Higgs boson contributions as signal		Higgs boson contributions as background	
	sequential-cut	BDT-based	sequential-cut	BDT-based
WZZ	5.2 ($3.7^{+2.2}_{-1.3}$)	6.1 ($3.8^{+2.2}_{-1.3}$)	5.8 ($3.7^{+2.3}_{-1.3}$)	5.8 ($3.7^{+2.3}_{-1.3}$)
ZZZ	5.4 ($6.0^{+4.6}_{-2.6}$)	5.4 ($6.2^{+4.9}_{-2.7}$)	5.6 ($6.3^{+5.3}_{-2.8}$)	5.7 ($6.3^{+5.3}_{-2.8}$)

Signal region	SS m_{jj} -in			SS m_{jj} -out			SS 1j			3ℓ		
	$e^\pm e^\pm$	$e^\pm \mu^\pm$	$\mu^\pm \mu^\pm$	$e^\pm e^\pm$	$e^\pm \mu^\pm$	$\mu^\pm \mu^\pm$	$e^\pm e^\pm$	$e^\pm \mu^\pm$	$\mu^\pm \mu^\pm$	0SFOS	1SFOS	2SFOS
Lost/three ℓ	1.4 ± 0.9	5.5 ± 1.6	7.0 ± 1.7	10.7 ± 2.6	9.7 ± 3.6	31.4 ± 3.8	2.5 ± 1.1	41.0 ± 6.1	5.8 ± 1.6	3.5 ± 0.7	25.6 ± 4.2	36.1 ± 3.1
Irreducible	1.0 ± 0.1	0.6 ± 0.1	2.9 ± 0.2	4.7 ± 0.4	1.9 ± 0.2	15.5 ± 1.2	0.4 ± 0.0	4.6 ± 0.2	0.5 ± 0.1	1.3 ± 0.1	1.2 ± 0.1	0.3 ± 0.0
Nonprompt ℓ	0.6 ± 0.6	3.6 ± 2.4	4.2 ± 1.5	0.8 ± 1.0	2.8 ± 1.5	9.1 ± 4.5	2.5 ± 5.2	2.9 ± 1.4	0.2 ± 0.1	1.8 ± 0.5	7.5 ± 2.3	1.8 ± 1.1
Charge flips	<0.1	<0.1	<0.1	4.5 ± 2.5	<0.1	<0.1	<0.1	0.1 ± 0.1	<0.1	<0.1	0.8 ± 1.2	0.3 ± 0.1
$\gamma \rightarrow$ nonprompt ℓ	0.1 ± 0.2	0.1 ± 0.4	<0.1	1.4 ± 0.5	1.1 ± 0.4	0.7 ± 0.4	0.6 ± 1.2	4.8 ± 8.0	<0.1	<0.1	1.0 ± 0.4	0.1 ± 1.5
Background sum	3.1 ± 1.1	9.8 ± 2.9	14.2 ± 2.3	22.1 ± 3.8	15.6 ± 4.0	56.8 ± 6.0	6.0 ± 5.4	53.5 ± 10.1	6.4 ± 1.6	6.6 ± 0.9	36.2 ± 5.0	38.7 ± 3.6
WWW onshell	0.9 ± 0.4	2.3 ± 0.9	4.6 ± 1.7	0.9 ± 0.4	1.0 ± 0.6	3.3 ± 1.3	0.3 ± 0.2	1.2 ± 0.4	0.4 ± 0.2	6.7 ± 2.4	4.3 ± 1.6	1.8 ± 0.7
WH \rightarrow WWW	0.4 ± 0.3	1.3 ± 0.9	1.2 ± 0.5	0.5 ± 0.3	1.3 ± 1.3	2.7 ± 1.2	1.1 ± 0.8	6.5 ± 3.1	2.2 ± 1.1	3.4 ± 1.6	5.0 ± 2.1	0.6 ± 0.6
WWW total	1.3 ± 0.5	3.7 ± 1.3	5.8 ± 1.7	1.5 ± 0.5	2.3 ± 1.4	6.0 ± 1.7	1.4 ± 0.8	7.7 ± 3.1	2.5 ± 1.1	10.1 ± 2.9	9.3 ± 2.6	2.4 ± 0.9
WWZ onshell	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	0.2 ± 0.1	<0.1	<0.1
ZH \rightarrow WWZ	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	0.1 ± 0.1	0.1 ± 0.1	<0.1
WWZ total	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	0.3 ± 0.1	0.1 ± 0.1	<0.1
WZZ onshell	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1
WH \rightarrow WZZ	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1
WZZ total	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1
ZZZ onshell	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1
ZH \rightarrow ZZZ	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1
ZZZ total	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1
VVV onshell	0.9 ± 0.4	2.3 ± 0.9	4.6 ± 1.7	0.9 ± 0.4	1.0 ± 0.6	3.3 ± 1.3	0.3 ± 0.2	1.2 ± 0.4	0.4 ± 0.2	6.9 ± 2.4	4.3 ± 1.6	1.8 ± 0.7
VH \rightarrow VVV	0.4 ± 0.3	1.3 ± 0.9	1.2 ± 0.5	0.5 ± 0.3	1.3 ± 1.3	2.7 ± 1.2	1.1 ± 0.8	6.5 ± 3.1	2.2 ± 1.1	3.6 ± 1.6	5.1 ± 2.1	0.6 ± 0.6
VVV total	1.3 ± 0.5	3.7 ± 1.3	5.8 ± 1.7	1.5 ± 0.5	2.3 ± 1.4	6.0 ± 1.7	1.4 ± 0.8	7.7 ± 3.1	2.5 ± 1.1	10.4 ± 2.9	9.3 ± 2.6	2.4 ± 0.9
Total	4.4 ± 1.2	13.5 ± 3.2	20.0 ± 2.9	23.6 ± 3.8	17.8 ± 4.2	62.7 ± 6.3	7.4 ± 5.5	61.2 ± 10.6	9.0 ± 2.0	17.0 ± 3.0	45.5 ± 5.6	41.1 ± 3.7
Observed	3	14	15	22	22	67	13	69	8	17	42	39


Signal region	$4\ell e\mu$					$4\ell ee/\mu\mu$		5ℓ	6ℓ
	bin 1	bin 2	bin 3	bin 4	bin 5	bin A	bin B		
ZZ	15.9±1.0	1.6±0.1	0.6±0.1	0.6±0.1	0.2±0.0	76.4±4.3	2.9±0.3	0.30±0.09	0.01±0.01
t \bar{t} Z	0.2±0.1	0.1±0.1	2.8±0.5	1.4±0.2	0.1±0.1	1.5±0.3	2.3±0.3	<0.01	<0.01
tWZ	0.1±0.1	0.1±0.1	0.6±0.1	0.7±0.1	0.1±0.1	0.5±0.1	0.7±0.1	<0.01	<0.01
WZ	0.5±0.2	0.2±0.2	0.5±0.2	0.3±0.3	0.1±0.1	1.0±0.4	0.2±0.1	<0.01	<0.01
Other	1.1±0.4	0.5±0.5	0.5±0.2	0.6±0.2	<0.1	2.7±0.6	0.5±0.2	<0.01	<0.01
Background sum	17.8±1.1	2.5±0.5	5.0±0.6	3.6±0.4	0.5±0.1	82.2±4.3	6.6±0.5	0.30±0.09	0.01±0.01
WWW onshell	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.01	<0.01
WH → WWW	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.01	<0.01
WWW total	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.01	<0.01
WWZ onshell	0.3±0.1	0.4±0.2	1.4±0.7	3.6±1.5	1.0±0.5	2.7±1.2	3.2±1.4	<0.01	<0.01
ZH → WWZ	1.1±0.5	1.1±0.5	0.5±0.2	1.3±0.5	1.8±0.8	2.9±1.2	1.5±0.6	<0.01	<0.01
WWZ total	1.3±0.5	1.5±0.5	1.9±0.8	4.9±1.6	2.9±0.9	5.6±1.7	4.7±1.5	<0.01	<0.01
WZZ onshell	0.2±0.2	0.1±0.1	0.2±0.2	0.4±0.4	0.1±0.1	0.5±0.4	0.2±0.2	2.62±1.82	0.03±0.05
WH → WZZ	0.2±0.3	0.2±0.3	<0.1	0.5±0.5	<0.1	<0.1	<0.1	<0.01	<0.01
WZZ total	0.4±0.3	0.3±0.3	0.2±0.2	0.9±0.7	0.1±0.1	0.5±0.4	0.2±0.2	2.62±1.82	0.03±0.05
ZZZ onshell	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.01	<0.01
ZH → ZZZ	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.01	<0.01
ZZZ total	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.01	<0.01
VVV onshell	0.5±0.2	0.4±0.2	1.6±0.8	4.0±1.5	1.1±0.5	3.2±1.3	3.4±1.4	2.62±1.82	0.03±0.05
VH → VVV	1.2±0.5	1.3±0.6	0.5±0.2	1.7±0.8	1.8±0.8	2.9±1.2	1.5±0.6	<0.01	<0.01
VVV total	1.7±0.6	1.7±0.6	2.1±0.8	5.8±1.7	3.0±0.9	6.1±1.8	4.8±1.5	2.62±1.82	0.03±0.05
Total	19.5±1.2	4.2±0.8	7.1±1.0	9.4±1.8	3.5±0.9	88.2±4.7	11.4±1.6	2.92±1.82	0.04±0.05
Observed	22	9	7	8	3	80	11	3	0

Signal region	SS m_{jj} -in			SS m_{jj} -out			SS 1j			3ℓ		
	$e^{\pm}e^{\pm}$	$e^{\pm}\mu^{\pm}$	$\mu^{\pm}\mu^{\pm}$	$e^{\pm}e^{\pm}$	$e^{\pm}\mu^{\pm}$	$\mu^{\pm}\mu^{\pm}$	$e^{\pm}e^{\pm}$	$e^{\pm}\mu^{\pm}$	$\mu^{\pm}\mu^{\pm}$	0 SFOS	1 SFOS	2 SFOS
Lost/three ℓ	1.8±0.4	10.9±2.0	8.7±1.0	8.8±1.7	46.0±6.2	44.8±4.4	8.4±1.3	43.5±4.4	34.5±2.7	4.6±0.8	15.1±1.5	58.3±2.4
Irreducible	2.1±0.4	13.0±3.6	8.4±1.4	9.8±1.4	41.1±4.5	42.8±4.7	2.6±0.6	22.8±8.6	13.2±1.9	2.5±0.9	2.2±1.2	2.5±0.8
Nonprompt ℓ	1.3±0.9	5.8±2.4	6.8±2.2	2.3±1.3	12.0±6.1	11.2±3.8	1.8±2.9	2.4±1.3	2.8±1.1	3.0±0.9	5.7±1.6	5.9±1.6
Charge flips	<0.1	1.2±2.0	<0.1	2.6±1.6	1.0±0.5	<0.1	6.9±4.7	0.2±0.1	<0.1	<0.1	1.1±1.3	0.7±0.2
$\gamma \rightarrow$ nonprompt ℓ	1.4±0.4	2.3±0.9	0.1±0.8	8.6±3.1	19.2±5.1	2.3±0.9	3.8±1.1	19.7±6.0	13.8±7.0	<0.1	0.6±0.7	0.2±0.3
Background sum	6.7±1.2	33.3±5.2	24.0±2.9	32.1±4.3	119±11	101±8	23.6±5.8	88.7±11.4	64.4±7.8	10.1±1.5	24.7±2.9	67.6±3.1
WWW onshell	1.0±0.5	3.3±1.5	3.5±1.6	0.9±0.5	3.9±1.8	4.1±1.9	0.5±0.3	1.8±0.8	1.7±0.9	5.9±2.6	3.8±1.7	2.5±1.2
WH \rightarrow WWW	0.2±0.3	1.9±1.5	0.6±0.4	0.4±0.4	1.3±0.8	1.7±1.0	0.8±0.5	4.5±2.7	3.3±2.0	3.0±1.7	2.7±1.5	1.3±0.8
WWW total	1.2±0.6	5.1±2.2	4.1±1.6	1.3±0.6	5.3±2.0	5.7±2.1	1.4±0.6	6.3±2.8	5.0±2.2	8.8±3.1	6.6±2.3	3.8±1.4
WWZ onshell	0.1±0.1	0.3±0.2	0.2±0.1	<0.1	<0.1	0.1±0.1	0.1±0.1	<0.1	<0.1	0.3±0.2	0.2±0.2	0.2±0.1
ZH \rightarrow WWZ	0.1±0.1	<0.1	<0.1	<0.1	<0.1	0.3±0.3	<0.1	<0.1	0.4±0.4	0.2±0.1	<0.1	<0.1
WWZ total	0.1±0.2	0.3±0.2	0.2±0.1	<0.1	<0.1	0.4±0.3	0.1±0.1	<0.1	0.4±0.4	0.4±0.2	0.2±0.2	0.2±0.1
WZZ onshell	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1
WH \rightarrow WZZ	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1
WZZ total	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1
ZZZ onshell	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1
ZH \rightarrow ZZZ	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1
ZZZ total	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1
VVV onshell	1.0±0.5	3.5±1.5	3.7±1.6	0.9±0.5	3.9±1.8	4.2±1.9	0.6±0.3	1.8±0.8	1.7±0.9	6.1±2.6	4.0±1.8	2.7±1.2
VH \rightarrow VVV	0.3±0.3	1.9±1.5	0.6±0.4	0.4±0.4	1.3±0.8	2.0±1.0	0.8±0.5	4.5±2.7	3.7±2.0	3.1±1.7	2.7±1.5	1.3±0.8
VVV total	1.3±0.6	5.4±2.2	4.2±1.6	1.3±0.6	5.3±2.0	6.1±2.1	1.4±0.6	6.3±2.8	5.4±2.2	9.3±3.1	6.8±2.3	3.9±1.4
Total	8.0±1.3	38.7±5.6	28.2±3.4	33.5±4.4	125±11	107±8	25.0±5.8	95.0±11.8	69.8±8.1	19.4±3.4	31.4±3.7	71.5±3.4
Observed	5	46	20	31	112	118	29	101	69	20	32	69

Signal region	$4\ell\ e\mu$				$4\ell\ ee/\mu\mu$			5ℓ	6ℓ
	bin 4	bin 3	bin 2	bin 1	bin A	bin B	bin C		
ZZ	0.3 ± 0.0	0.7 ± 0.0	0.7 ± 0.0	0.4 ± 0.0	1.8 ± 0.2	6.0 ± 0.6	5.0 ± 0.5	0.30 ± 0.08	0.01 ± 0.01
$t\bar{t}Z$	0.2 ± 0.0	0.3 ± 0.1	0.8 ± 0.1	2.3 ± 0.4	1.4 ± 0.2	1.1 ± 0.2	0.2 ± 0.0	<0.01	<0.01
tWZ	0.1 ± 0.1	0.1 ± 0.1	0.3 ± 0.0	0.8 ± 0.1	0.5 ± 0.1	0.3 ± 0.1	0.1 ± 0.1	<0.01	<0.01
WZ	0.2 ± 0.1	0.1 ± 0.1	0.1 ± 0.2	0.6 ± 0.2	<0.1	0.2 ± 0.1	0.1 ± 0.1	<0.01	<0.01
Other	<0.1	0.2 ± 0.1	0.6 ± 0.3	0.2 ± 0.1	<0.1	1.4 ± 0.5	0.1 ± 0.1	<0.01	<0.01
Background sum	0.8 ± 0.1	1.4 ± 0.1	2.5 ± 0.3	4.3 ± 0.4	3.7 ± 1.9	9.1 ± 0.8	5.5 ± 0.5	0.30 ± 0.08	0.01 ± 0.01
WWW onshell	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.01	<0.01
$WH \rightarrow WWW$	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.01	<0.01
WWW total	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.01	<0.01
WWZ onshell	0.5 ± 0.2	0.5 ± 0.2	1.1 ± 0.4	4.0 ± 1.6	2.1 ± 0.9	1.2 ± 0.4	0.6 ± 0.2	<0.01	<0.01
$ZH \rightarrow WWZ$	2.3 ± 0.9	1.1 ± 0.4	0.3 ± 0.1	0.1 ± 0.1	0.8 ± 0.3	0.9 ± 0.4	0.5 ± 0.2	<0.01	<0.01
WWZ total	2.8 ± 0.9	1.6 ± 0.5	1.4 ± 0.4	4.1 ± 1.6	2.9 ± 1.0	2.1 ± 0.6	1.1 ± 0.3	<0.01	<0.01
WZZ onshell	<0.1	0.1 ± 0.1	0.1 ± 0.1	0.4 ± 0.3	0.2 ± 0.2	0.1 ± 0.1	0.1 ± 0.1	2.17 ± 1.46	0.03 ± 0.04
$WH \rightarrow WZZ$	<0.1	0.4 ± 0.3	0.1 ± 0.2	<0.1	<0.1	<0.1	<0.1	<0.01	<0.01
WZZ total	<0.1	0.4 ± 0.4	0.2 ± 0.2	0.4 ± 0.3	0.2 ± 0.2	0.1 ± 0.1	0.1 ± 0.1	2.17 ± 1.46	0.03 ± 0.04
ZZZ onshell	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.01	<0.01
$ZH \rightarrow ZZZ$	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.01	<0.01
ZZZ total	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.01	<0.01
VVV onshell	0.5 ± 0.2	0.6 ± 0.2	1.2 ± 0.4	4.4 ± 1.6	2.3 ± 0.9	1.3 ± 0.5	0.7 ± 0.2	2.17 ± 1.46	0.03 ± 0.04
$VH \rightarrow VVV$	2.3 ± 0.9	1.5 ± 0.5	0.4 ± 0.3	0.1 ± 0.1	0.8 ± 0.3	0.9 ± 0.4	0.5 ± 0.2	<0.01	<0.01
VVV total	2.8 ± 0.9	2.1 ± 0.6	1.6 ± 0.5	4.5 ± 1.6	3.1 ± 1.0	2.2 ± 0.6	1.2 ± 0.3	2.17 ± 1.46	0.03 ± 0.04
Total	3.6 ± 0.9	3.5 ± 0.6	4.1 ± 0.6	8.8 ± 1.7	6.8 ± 2.1	11.3 ± 1.0	6.6 ± 0.6	2.47 ± 1.46	0.04 ± 0.04
Observed	7	1	5	7	6	8	7	3	0


History lesson

m_{top} vs. m_W and m_{Higgs}




...after analysis of Run I data, ... **②** m_W shifted a full s.d. ... the m_{Higgs} must be **③** much lower than anyone had anticipated. ... Surprises happen.

— D. Amidei, R. Brock Fermi news 1/17/2003




History tells us with more data we get smarter; also surprises happen




ment with the results obtained from simulation. The $\sigma(p_T)/p_T$ averaged over ϕ and η varies in p_T from $(1.8 \pm 0.3(\text{stat.}))\%$ at $p_T = 30 \text{ GeV}/c$ to $(2.3 \pm 0.3(\text{stat.}))\%$ at $p_T = 50 \text{ GeV}/c$, again in good agreement with the expectations from simulation.

<https://arxiv.org/pdf/1206.4071.pdf>



(a)



(b)

arXiv.org > physics > arXiv:1502.02701

Search...

Help | Advanced


Physics > Instrumentation and Detectors

[Submitted on 9 Feb 2015 (v1), last revised 1 Jul 2015 (this version, v2)]

Performance of electron reconstruction and selection with the CMS detector in proton–proton collisions at $\sqrt{s} = 8$ TeV

CMS Collaboration

The performance and strategies used in electron reconstruction and selection at CMS are presented based on data corresponding to an integrated luminosity of 19.7 inverse femtobarns, collected in proton–proton collisions at $\sqrt{s} = 8$ TeV at the CERN LHC. The paper focuses on prompt isolated electrons with transverse momenta ranging from about 5 to a few 100 GeV. A detailed description is given of the algorithms used to cluster energy in the electromagnetic calorimeter and to reconstruct electron trajectories in the tracker. The electron momentum is estimated by combining the energy measurement in the calorimeter with the momentum measurement in the tracker. Benchmark selection criteria are presented, and their performances assessed using Z, Upsilon, and J/psi decays into electron–positron pairs. The spectra of the observables relevant to electron reconstruction and selection as well as their global efficiencies are well reproduced by Monte Carlo simulations. The momentum scale is calibrated with an uncertainty smaller than 0.3%. The momentum resolution for electrons produced in Z boson decays ranges from 1.7 to 4.5%, depending on electron pseudorapidity and energy loss through bremsstrahlung in the detector material.



<https://twiki.cern.ch/twiki/pub/CMSPublic/BTV13TeV2017FIRST2018/PT30GeV.pdf>

